


IL27

Mouse Anti-Human IL-27 (Clone B-G49) FITC mAb

Catalog No.	CDM284	Quantity:	100 tests
Alternate Names:	Interleukin 27; p28; IL30; IL27A; IL-27A; IL27p28		
Description:	The monoclonal antibody recognizes human IL-27, one of the subunits of a heterodimeric cytokine complex. which interacts with Epstein-Barr virus induced gene 3 (EBI3), and forms a complex that has been shown to drive rapid expansion of naive but not memory CD4(+) T cells. The complex is also found to synergize strongly with IL-12 to trigger IFN- γ production of naive CD4(+) T cells.		
Concentration:	100 tests/ 1 ml		
Gene ID:	246778		
Conjugate	FITC		
Specificity:	Recognizes both natural and recombinant human IL-27.		
Host:	Mouse		
Isotype:	IgG2a		
Immunogen:	Recombinant human IL-27		
Clone:	B-G49		
Formulation:	Liquid in PBS with 5 % BSA and 0.1 % sodium azide. Precaution: Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.		
Purification:	Ion exchange chromatography		
Applications:	Flow Cytometry		
Application Notes:	Use 10 μ l to label 10^6 cells or 100 μ l of whole blood. The optimal concentration should be determined by the user for each specific application.		
Storage & Stability:	Stable at 2-8°C for 1 year. For longer storage, freeze in aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		

IL-27 FITC staining of cultured dendritic cells:(left) unactivated and (right) activated with 1 μ g/ml LPS, 10ng/ml IFN-g and 100ng/ml TNF-a, 3×10^5 dendritic cells were labelled with 10 μ L of anti-IL-27 FITC (green) or a FITC conjugated IgG2a isotype control (black).



cellsciences.com

10³ 10⁴ 10⁵ FL1-H 10³ 10⁴ 10⁵ FL1-H 10³ 10⁴ 10⁵ FL1-H 10³ 10⁴ 10⁵

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com