

CALR

Rabbit Anti-Human Calreticulin pAb

Catalog No.	CPC134	Quantity:	100 µg
Alternate Names:	CRT, FLJ26680, RO, SSA, cC1qR, Sicca syndrome antigen A (autoantigen Ro; calreticulin), autoantigen Ro		
Description:	Rabbit Anti-Human Calreticulin polyclonal antibody. Calreticulin is a calcium binding protein found in abundance in the endoplasmic reticulum and the sarcoplasmic reticulum. Like many other ER proteins, it has the conserved ER retention KDEL (Lys-Asp-Glu-Leu) sequence at its C-terminus. Calreticulin has also been detected in the nucleus and nuclear envelop. Recent studies suggest that this soluble ER protein has a multifunctional role as it appears to be involved in calcium storage and regulation as well as having a molecular chaperone activity. Studies also suggest its involvement in certain autoimmune diseases. Consistent with its multiple functions, the calreticulin molecule appears to have a zonal character. The protein has both high and low affinity calcium binding sites.		
Concentration:	0.2 mg/ml		
Gene ID:	811		
Specificity:	Recognizes a 60 kDa calreticulin on SDS-PAGE immunoblots.		
Host:	Rabbit		
Immunogen:	Synthetic peptide mapping to the C-terminus of human calreticulin		
Isotype:	IgG		
Formulation:	Liquid in PBS + 30% glycerol + 0.5 mg/ml BSA + 0.01% thimerosal. Precaution: Thimerosal is a poisonous and hazardous substance which should be handled by trained staff only.		
Purification:	Affinity purified		
Reconstitution:	The antibody solution should be gently mixed before use.		
Cross-Reactivity:	Human, mouse, and rat		
Applications:	Western Blot Immunohistochemistry		
Application Notes:	For Western Blot, use a working dilution of 0.5-4 µg/ml. For Immunoprecipitation, use a working dilution of 10-20 µg/ml. The optimal concentration should be determined by the user for each specific application.		
Storage & Stability:	Store at -20°C or in working aliquots at -80°C for long term storage. Avoid repeated freeze-thaw cycles.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

