

CCL14

Recombinant Human Chemokine (CC motif) Ligand 14

Catalog No.	CRH100A CRH100B CRH100C	Quantity:	2 µg 10 µg 1.0 mg
Alternate Names:	HCC-1, CC-1, CC-3, CKb1, HCC-3, MCIF, NCC2, SCYA14, SCTL2, SY14		
Gene ID:	6358		
Protein Accession No:	Q16627		
Description:	<p>Recombinant Human CCL14 is a single, non-glycosylated polypeptide chain containing 72 amino acids.</p> <p>Background: HCC-1/CCL14 is a CC chemokine that signals through the CCR1 receptor and chemoattracts blood monocytes. It is secreted by various tissues including skeletal muscle, heart, spleen, liver and bone marrow.</p> <p>It is a single non-glycosylated polypeptide chain containing 72 amino acids.</p>		
Source:	<i>E. coli</i>		
Molecular Weight:	8.4 kDa		
Formulation:	Lyophilized from a sterile filtered solution containing 20 mM PBS, pH 7.4 + 100 mM NaCl		
Purity:	> 96% as determined by HPLC and SDS-PAGE analyses		
Endotoxin Level:	Less than 1EU/µg as determined by LAL method.		
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ determined by a chemotaxis bioassay using human monocytes is in a concentration of 5.0-20 ng/ml.		
Amino Acid Sequence:	TESSSRGPYH PSECCFTYTT YKIPRQRIMD YYETNSQCSK PGIVFITKRG HSVCTNPSDK WVQDYIKDMK EN		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water or aqueous buffer to a concentration of 0.1-1.0 mg/ml. Further dilutions should be made in appropriate buffered solutions.		
Storage & Stability:	This lyophilized preparation is stable at 2-4°C, but should be kept desiccated at -20°C for long term storage. Upon reconstitution, the preparation is stable for up to one week at 2-4°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -80°C. Avoid repeated freeze/thaw cycles.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.