

## Sonic Hedgehog, Recombinant Human Mature Peptide (Amino Terminal Peptide), Carrier-Free

**Catalog No:** CRS600A    **Quantity:** 5 µg  
CRS600B                    25 µg  
CRS600C                    1.0 mg

**Description:** The *hedgehog* (hh) gene encoding a secreted protein was originally identified in *Drosophila* as a segment polarity gene. The vertebrate homologues of Hh comprise several proteins including sonic hedgehog (Shh), Indian hedgehog (Ihh), and Desert hedgehog (Dhh). Hedgehog proteins are important signaling molecules during embryonic development. Shh genes are highly conserved and have been identified in a variety of species including human, mouse, frog, fish, and chicken. Mouse and human Shh are 92% identical at the amino acid sequence level. Shh is expressed in key embryonic tissues such as the Hensen's node, the zone of polarizing activity in the posterior limb bud, the notochord, and the floor plate of the neural tube. Shh is involved in regulating the patterning of the developing central nervous system, somite, and limb. Shh plays an important role in the development of particular tissues such as whisker, hair, foregut, tooth and bone. Recent evidence also suggests that Shh is involved in regulating stem cell fates of neural and hematopoietic lineages, and that aberrant Shh signaling is implicated in basal cell carcinomas and other diseases.

Human Shh cDNA encodes a 45 kDa precursor protein. An autocatalytic reaction yields a 19 kDa amino-terminal domain Shh-N protein containing cholesterol and palmitate, and a 25 kDa carboxy-terminal domain Shh-C protein. The N-terminal domain retains all known signaling capabilities, while the C-terminal domain is responsible for the intramolecular processing, acting as a cholesterol transferase. Shh can act as both a short-range contact dependent factor and as a long-range, diffusible morphogen. At the cell surface, Shh activity is mediated by a multicomponent receptor complex involving the 12-pass transmembrane protein Patched (Ptc) which binds Shh with high affinity and Smoothed (Smo), a signaling seven transmembrane G-protein coupled receptor. In the absence of Shh, Ptc represses Smo activity. The binding of Shh to Ptc, releases the basal repression of Smo by Ptc.<sup>1-5</sup>

**Source:** A DNA sequence encoding amino acid residues Cys 24-Gly 197 of Human Sonic Hedgehog was fused to a 6X histidine tag at the carboxy-terminus. The fusion protein was expressed in *E. coli*.

**Purity:** Greater than 97% as determined by SDS-PAGE and visualized by silver stain.

**Endotoxin Level:** Less than 1.0 EU per 1 µg of the cytokine as determined by the LAL method.

**Formulation:** Supplied as a 0.2 µm filtered solution (85.9 µL) in PBS at a concentration of 0.294 mg/ml.

**Stability:** This cytokine, in the presence of a carrier protein, can be stored under sterile condition at 2° - 8°C for one month or at -20°C to -70°C in a **manual defrost freezer** for three months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**



**Cell Sciences<sup>TM</sup>, Inc.**  
Neponset Valley Tech Park  
480 Neponset St., Bldg. 12A  
Canton, MA 02021

Toll Free: 888 769-1246  
Phone: 781 828-0610  
Fax: 781 828-0542

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Web Site: [www.cellsciences.com](http://www.cellsciences.com)