

BRCA1


Rabbit Anti-Human BReast CAncer 1 early onset Clone Poly6121 pAb

Catalog No.	CSI14349 CSI14350	Quantity:	50 µl 200 µl
Alternate Names:	Bcl-2, B-cell lymphoma protein 2, Breast and ovarian cancer susceptibility protein 1		
Description:	<p>BRCA1 (breast and ovarian cancer susceptibility protein 1) is a RING finger protein containing a BRCT domain. BRCA1 exists as a heterodimer with 22 possible isoforms. The full length protein has a reported molecular weight of 208 kD. BRCA1 localizes to the mitotic spindle microtubules, centriole walls, pericentriolar fibers at centrosomes. Unphosphorylated BRCA1 localizes on chromosomes from metaphase through telophase; phosphorylated BRCA1 resides in inner chromosomal structure, centrosome, cleavage furrow during prophase through telophase, and relocates to the perinuclear region when cells are subjected to IR or UV radiation in S phase. BRCA1 acts as a tumor suppressor and can function as a secreted growth inhibitory protein, participates in transcription-coupled repair of oxidative DNA damage, and X-chromosome inactivation. BRCA1 can also function as a E3 ubiquitin ligase. BRCA1 can be transcriptionally downregulated by Ets-2, Brg-1, and Hmga-1. BRCA1 can be modified by glycosylation, ubiquitination and phosphorylation by CDK4, ATM/ATR, cdk2, and hChk2. The BRCA1 protein has been reported to interact with RNA polymerase II holoenzyme and BARD1. The Poly6121 antibody has been shown to be useful for Western blotting of the human BRCA1 protein.</p>		
Structure:	RING finger, BRCT domain, heterodimer. 22 possible isoforms, full length 208 kD.		
Gene ID:	672		
Distribution:	<p>Mitotic spindle microtubules, centriole walls, pericentriolar fibers at centrosomes. Unphosphorylated BRCA1 localizes on chromosomes from metaphase through telophase. Phosphorylated BRCA1 resides in inner chromosomal structure, centrosome, cleavage furrow during prophase through telophase, and relocates to the perinuclear region when cells are subjected to IR or UV radiation in S phase.</p>		
Function:	Tumor suppressor. Secreted growth inhibitory protein. Involved in transcription-coupled repair of oxidative DNA damage, X-chromosome inactivation, E3 ubiquitin ligase activity.		
Host:	Rabbit		
Immunogen:	Recombinant (partial), N-terminal		
Isotype:	IgG		
Clone:	Poly6121		



- Regulation:** Transcriptionally downregulated by Ets-2, Brg-1, Hmga-1. Phosphorylation by CDK4, ATM/ATR, cdk2, and hChk2
- Formulation:** This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 50% glycerol. **Precaution:** Sodium azide is a poisonous and hazardous substance which should be handled by trained staff only.
- Purification:** The antibody was purified by antigen-affinity chromatography.
- Modification:** Phosphorylation, Glycosylation, Ubiquitination
- Reactivity:** Human
- Applications:** Western Blot
- Recommended Usage:** Each lot of this antibody is quality control tested by Western blotting. Western blotting, suggested working dilution(s): Use 10 µl per 5 ml antibody dilution buffer for each mini-gel. It is recommended that the reagent be titrated for optimal performance for each application
- Storage & Stability:** Upon receipt, store frozen at -20° C.
- Interaction:** RNA polymerase II holoenzyme, BARD1

A431 cell extract was resolved by electrophoresis, transferred to nitrocellulose and probed with rabbit anti-BRCA1 antibody. Proteins were visualized using a donkey anti-rabbit secondary conjugated to HRP and a chemiluminescence detection system.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

