

IGF-II, Human pAb

Catalog No.	PAC1 5PAC1	Quantity:	20 µl (~500 RIA tubes) 100 µl (~2,500 RIA tubes)
Specificity:	Antibody reacts 100% with human IGF-II, rat IGF-II, chicken IGF-II and pig IGF-II. Reacts 85% bovine IGF-II.		
Host:	Rabbit		
Immunogen:	Recombinant human IGF-II		
Cross-Reactivity:	Less than 0.1% cross-reactive with ovine insulin, human IGF-I, chicken IGF-I, and rat IGF-I.		
Applications:	Radioimmunoassay: working dilution of 1:7,500 At a dilution of 1:80,000 the antiserum binds 20% of fresh [¹²⁵ I]hIGF-II tracer (80 Ci/g). Half-maximal inhibition (ED ₅₀) of radioligand binding is produced by ~ 4 ng/ml of hIGF-II. The optimal concentration should be determined by the user for each specific application.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Web site: www.cellsciences.com