

## Recombinant Human BCL2 minus BH1 domain His

<b>Catalog No.</b>	CRB129A	<b>Quantity:</b>	2 µg
	CRB129B		10 µg
	CRB129C		1.0 mg

**Alternate Names:** B-cell CLL/lymphoma-2, BCL-2, PCKBCL2

**Description:** BCL2 gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends.  
Bcl-2 Des BH1 domain (136-155 residues) Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 198 amino acids 1-135 and 156-218. The Bcl-2 is expressed as His-Tag fusion protein and purified by proprietary chromatographic techniques.

**Gene ID:** 596

**Source:** *E. coli*

**Formulation:** Lyophilized from a sterile filtered solution containing 10 mM Tris-HCl, pH 8.0 + 1 mM EDTA + 250 mM NaCl

**Purity:** > 95% as determined by RP-HPLC and SDS-PAGE analyses

**Endotoxin Level:** < 0.1 ng/µg of BCL2

**Reconstitution:** **Centrifuge vial prior to opening.** Suspend BCL2 in 100 µl of 0.5 M Acetic acid, overnight at 2-4°C. Dilute 10 fold into selected buffer system. Because BCL2 has a tendency to form intramolecular disulfide bond, 5 mM DTT is recommended in assay buffer. When running SDS-PAGE gel, 10 mM DTT is recommended.

Input marker or positive control (Western Blot)

Function study (Binding Assay)

The optimal concentration should be determined by the user for each specific application.

**Storage & Stability:** Lyophilized Bcl-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bcl-2 should be stored at 4°C between 2-7 days and for future use below -18°C.  
For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).  
**Please note that the addition of any carrier protein into this product may produce unwanted endotoxin. This depends upon the particular application employed.**  
**Avoid repeated freeze-thaw cycles.**

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



**Cell Sciences®**  
480 Neponset Street  
Bldg 12A  
Canton, MA 02021

Toll Free: 888-769-1246  
Phone: 781-828-0610  
Fax: 781-828-0542

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)