


## WISP2

### Recombinant Human WISP-2 / CTGF-L

<b>Catalog No.</b>	CRC603A CRC603B CRC603C	<b>Quantity:</b>	5 µg 20 µg 1.0 mg
<b>Alternate Names:</b>	WNT1-inducible-signaling pathway protein 2, Connective tissue growth factor-like protein, CTGF-L, CCN family member 5		
<b>Description:</b>	WNT1-inducible-signaling pathway protein 2 (WISP-2) is a member of the CYR61/CTGF/NOV (CCN) family of regulatory factors. WISP-2 is expressed in ectodermal, mesodermal, and endodermal lineages, including primary osteoblasts, fibroblasts, mesenchymal stem cells, and adipogenic precursor cells. WISP-2 is a canonical WNT ligand that regulates cell proliferation, adhesion, and metastasis. Secreted WISP-2 promotes mesenchymal precursor cell proliferation and maintains them in an undifferentiated state. In bone-forming osteoblasts, WISP-2 promotes osteoblast adhesion and inhibits osteocalcin production.		
<b>UniProt ID:</b>	O76076		
<b>Gene ID:</b>	8839		
<b>Source:</b>	<i>E. coli</i>		
<b>Molecular Weight:</b>	24.8 kDa (228 aa) monomer		
<b>Formulation:</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)		
<b>Purity:</b>	≥ 95% by reducing and non-reducing SDS-PAGE		
<b>Endotoxin Level:</b>	< 1 EU/µg by kinetic LAL		
<b>Amino Acid Sequence:</b>	MQLCPTPCTC PWPPPRCPLG VPLVLDGCGC CRVCARRLGE PCDQLHVCD SQGLVCQPGA GPGGRGALCL LAEDDSSCEV NGRLYREGET FQPHCSIRCR CEDGGFTCVPLCSEDVRLPS WDCPHPRRVE VLKCCPEWV CGQGGGLGTQ PLPAQGPQFS GLVSSLPPGV PCPEWSTAWG PCSTTCGLGM ATRVSNQNR FCRLETQRRLC LSRPCPPSRG RSPQNSAF		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipet the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for reconstitution. A small amount of precipitate may be seen.		
<b>Storage &amp; Stability:</b>	Store as supplied at -20°C to -80°C for up to 1 year. Upon reconstitution, prepare working aliquots and store at -20°C to -80°C. It is recommended that a carrier protein such as 0.1% HSA or BSA is added for long term storage. <b>Avoid repeated freeze-thaw cycles.</b>		





**Human WISP-2 / CTGF-L Gel**

Figure: 1 ug run under (-) non-reducing and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human WISP-2 is predicted to have a MW of 24.4 kDa.

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)