

FLT1

Recombinant Human VEGFR-1 (D4), soluble

Catalog No.	CRF111A CRF111B CRF111C	Quantity:	5 µg 20 µg 1 mg
Alternate Names:	Human Vascular Endothelial Growth Factor Receptor 1, Fms-related Tyrosine Kinase 1, FLT-1		
Description:	<p>Recombinant Human soluble Vascular Endothelial Growth Factor Receptor-1 domain D1-4 is produced as a non-chimeric protein in a monomeric form. The soluble receptor protein contains only the first 4 extracellular domains, which contain all the information necessary for binding of VEGF. Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVEC supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA. The biological functions of sVEGFR-1 still are not clear, but it seems to be an endogenous regulator of angiogenesis, binding VEGF with the same affinity as the full-length receptor.</p>		
UniProt ID:	P17948		
Gene ID:	2321		
Source:	Insect cells		
Molecular Weight:	55 kDa (457 aa), monomer		
Formulation:	Lyophilized from PBS		
Purity:	> 90%, by SDS-PAGE and visualized by silver stain		
Endotoxin Level:	< 1.0 EU/µg		
Biological Activity:	The activity of sVEGFR-1(D4) was determined by its ability to inhibit the VEGF-A-induced proliferation of HUVECs.		
Reconstitution:	Centrifuge vial prior to opening. Soluble in water and most aqueous buffers. Add sterile distilled water to the vial to fully solubilize the protein to a concentration of ≥ 0.10 mg/ml.		
Storage & Stability:	Stable for up to 1 year at -20°C to -80°C or for 6 months in working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		

Amino Acid Sequence: SKLKDPESLSLKGQTQHIMQAGQTLHLQCRGEEAAHKWSLPEMVSKESERLSITKSACGRN
GKQFCSTLTLNTAQANHTGFYSCKYLAVPTSKKKETESAIYIFISDTGRPFVEMYSEIPEII
HMTEGRELVIPCRVTSPNITVTLKKFPLDTLIPDGKRIIWDSRKGFIISNATYKEIGLLTCEA
TVNGHLYKTNYLTHRQNTIIDVQISTPRPVKLLRGHTLVLNCTATTPLNTRVQMTWSYP
DEKNKRASVRRRIDQSNSHANIFYSVLTIDKMQNKDKGLYTCRVRSGPSFKSVNTSVHI
YDKAFITVKHRKQQVLETVAGKRSYRLSMKVKAFFSPEVVWLKDGLPATEKSARYLTRG
YSLIHKDVTEEDAGNYTILLSIKQSNVFKNLTATLIVNVKPKIYEKAVSSFPDPALYPLGSR
QILTCTAYGIPQPTIKWFWHPCNHNHSEARCDFC

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com