

ACVR1


Recombinant Human ALK-2 / ACVR1 (His Tag)

Catalog No.	CRH419A-His CRH419B-His	Quantity:	100 µg 200 µg
Alternate Names:	Activin receptor type-1, Activin receptor type I, ACTR-I, Activin receptor-like kinase 2, ALK-2, Serine/threonine-protein kinase receptor R1, SKR1, TGF-B superfamily receptor type I, TSR-I		
Description:	<p>ALK-2, also termed as ACVR1, was initially identified as an activin type I receptor because of its ability to bind activin in concert with ActRII or ActRIIB. ALK-2 is also identified as a BMP type I receptor. It has been demonstrated that ALK-2 forms complex with either the BMP-2/7-bound BMPRII or ACVR2A /ACVR2B. ALK-1 and ALK-2 presenting in the yeast <i>Saccharomyces cerevisiae</i> are two haspin homologues. Both ALK-1 and ALK-2 exhibit a weak auto-kinase activity in vitro, and are phosphoproteins in vivo. ALK-1 and ALK-2 levels peak in mitosis and late-S/G2. Control of protein stability plays a major role in ALK-2 regulation. The half-life of ALK-2 is particularly short in G1. Overexpression of ALK-2, but not of ALK-1, causes a mitotic arrest, which is correlated to the kinase activity of the protein. This suggests a role for ALK-2 in the control of mitosis. Endoglin is phosphorylated on cytosolic domain threonine residues by the TGF-beta type I receptors ALK-2 and ALK-5 in prostate cancer cells. Endoglin did not inhibit cell migration in the presence of constitutively active ALK-2. Defects in ALK-2 are a cause of fibrodysplasia ossificans progressiva (FOP).</p>		
UniProt ID:	Q04771		
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-Val 124) of human ALK2 (Met 1-Val 124) was fused with a polyhistidine tag at the C-terminus.		
Source:	Baculovirus-Insect Cells		
Formulation:	Supplied as sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% gly Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted recombinant human ALK2 consists of 114 amino acids and has a calculated molecular mass of 12.8KDa. It migrates as an approximately 17 kDa band in SDS-PAGE under reducing conditions.		
Purity:	> 93 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Kinase activity untested		
Predicted N-terminal:	Met 21		
Reconstitution:	<p>Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.</p>		

Storage & Stability:

Stable for up to 1 year from date of receipt at -20°C to -80°C
After reconstitution, store working aliquots at -20°C to -80°C .

Avoid repeated freeze-thaw cycles.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.