

ACVR2A



Recombinant Human ACVR2 / ACTRII / ACVR2A (His Tag)

Catalog No.	CRH428A-His CRH428B-His	Quantity:	100 µg 200 µg
Alternate Names:	Activin receptor type-2A, Activin receptor type IIA, ACTR-IIA, ACTRIIA		
Description:	ACVR2A and ACVR2B are two activin type II receptors. ACVR2A has been shown to interact with INHBA, SYNJ2BP and ACVR1B. The bovine ACVR2A gene encodes a protein of 513 amino acids which is highly homologous (approximately 98% identity) to the rat, mouse, and human ACVR2A proteins. Inactivation of ACVR2A is a common event in prostate cancer cells suggesting it may play an important role in the development of prostate cancer. The ACVR2A gene is a putative tumor suppressor gene that is frequently mutated in microsatellite-unstable colon cancers (MSI-H colon cancers). Frameshift mutation of ACVR2A may contribute to MSI-H colon tumorigenesis via disruption of alternate TGF-beta effector pathways.		
UniProt ID:	P27037		
Accession Number:	NP_001607.1		
Protein Construction:	A DNA sequence encoding the extracellular domain of human ACVR2A (Met 1-Pro 134) was expressed, with a polyhistidine tag at the C-terminus.		
Source:	HEK293 Cells		
Molecular Weight:	The rhACVR2A comprises 126 amino acids and predicts a molecular mass of 14.9 kDa. As a result of glycosylation, the apparent molecular mass of the protein is ~35-40 kDa in SDS-PAGE under reducing conditions.		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Purity:	> 97 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED50 for this effect is typically 0.4-3 µg/mL in the presence of 10 ng/mL recombinant Activin A.		
Predicted N-terminal:	Ala 20		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		



Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED50 for this effect is typically 0.4-3 $\mu\text{g}/\text{mL}$ in the presence of 10 ng/mL recombinant Activin A.

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com