

BMPR1A


Recombinant Human BMPR-1A / ALK-3 / CD292 (His Tag)

Catalog No.	CRH484A-His CRH484B-His	Quantity:	100 µg 200 µg
Alternate Names:	Bone morphogenetic protein receptor type-1A, BMP type-1A receptor, BMPR-1A, Activin receptor-like kinase 3, ALK-3, Serine/threonine-protein kinase receptor R5, SKR5, CD292		
Description:	<p>Bone Morphogenetic Protein Receptor, type IA (BMPR-1A), is a type I receptor for bone morphogenetic proteins (BMPs) which belong to the transforming growth factor beta (TGF-β) superfamily. The BMP receptors form a subfamily of transmembrane serine/threonine kinases including the type I receptors BMPR-1A and BMPR-1B and the type II receptor BMPR2. BMPR-1A is expressed in the epithelium during branching morphogenesis. Deletion of BMPR-1A in the epithelium with an Sftpc-cre transgene leads to dramatic defects in lung development. ALK-3 and ALK-6 share a high degree of homology, yet possess distinct signaling roles. The transforming growth factor (TGF)-beta type III receptor (TbetaRIII) enhanced both ALK-3 and ALK-6 signaling. TbetaRIII associated with ALK-3 primarily through their extracellular domains, whereas its interaction with ALK-6 required both the extracellular and cytoplasmic domains. ALK-3 plays an essential role in the formation of embryonic ventral abdominal wall, and abrogation of BMP signaling activity due to gene mutations in its signaling components could be one of the underlying causes of omphalocele at birth. The type IA BMP receptor, ALK-3 was specifically required at mid-gestation for normal development of the trabeculae, compact myocardium, interventricular septum, and endocardial cushion. Cardiac muscle lacking ALK-3 was specifically deficient in expressing TGFbeta2, an established paracrine mediator of cushion morphogenesis.</p>		
UniProt ID:	P36894		
Accession Number:	NP_004320.2		
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-Arg 152) of human ALK3 was fused with a polyhistidine tag at the C-terminus.		
Source:	Hek293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted recombinant human ALK3 consists of 141 amino acids and has a predicted molecular mass of 15.7 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rhALK3 is approximately 28-33 kDa due to glycosylation.		
Purity:	> 85 % as determined by SDS-PAGE		
Endotoxin Level:	< 1.0 EU per µg protein as determined by the LAL method.		




- Biological Activity:** Measured by its ability to inhibit BMP4-induced alkaline phosphatase production by MC3T3E1 mouse preosteoblast cells. The ED50 for this effect is 20-120 ng/mL.
- Predicted N-terminal:** Gln 24
- Reconstitution:** **Centrifuge vial prior to opening.** Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. **DO NOT VORTEX.** Allow several minutes for complete reconstitution.
- Storage & Stability:** Stable for up to 1 year from date of receipt at -20°C to -80°C. After reconstitution, store working aliquots at -20°C to -80°C. **Avoid repeated freeze-thaw cycles.**

Measured by its ability to inhibit BMP4-induced alkaline phosphatase production by MC3T3E1 mouse preosteoblast cells. The ED50 for this effect is 2.5-10 µg/mL.



SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com