


## Angptl4

### Recombinant Mouse Angiopoietin-related protein 4 (His Tag)

<b>Catalog No.</b>	CRM596A-His CRM596B-His	<b>Quantity:</b>	50 µg 100 µg
<b>Alternate Names:</b>	Angiopoietin-related protein 4, 425O18-1, Angiopoietin-like protein 4, Fasting-induced adipose factor, Hepatic fibrinogen/angiopoietin-related protein, HFARP, Secreted protein Bk89		
<b>Description:</b>	ANGPTL4, also known as ANGPTL2, is a protein with hypoxia-induced expression in endothelial cells. It contains 1 fibrinogen C-terminal domain and is expressed at high levels in the placenta, heart, liver, muscle, pancreas and lung but expressed poorly in the brain and kidney. ANGPTL4 inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage. It may act as a regulator of angiogenesis and modulate tumorigenesis. It inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage. It may also exert a protective function on endothelial cells through an endocrine action. ANGPTL4 is directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity. In response to hypoxia, the unprocessed form of the protein accumulates in the subendothelial extracellular matrix (ECM). The matrix-associated and immobilized unprocessed form limits the formation of actin stress fibers and focal contacts in the adhering endothelial cells to inhibit their adhesion and decreases their motility.		
<b>UniProt ID:</b>	Q9Z1P8		
<b>Protein Construction:</b>	A DNA sequence encoding the mouse ANGPTL4 (Lys167-Ser410) was expressed with a polyhistidine tag at the N-terminus.		
<b>Source:</b>	HEK293 Cells		
<b>Formulation:</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
<b>Molecular Weight:</b>	The rmANGPTL4 consists of 264 aa with a predicted MW of 29.9 KDa and migrates at ~38-45 KDa in SDS-PAGE under reducing conditions, due to glycosylation.		
<b>Purity:</b>	> 95 % as determined by SDS-PAGE.		
<b>Endotoxin Level:</b>	< 1.0 EU per µg of the protein as determined by the LAL method		
<b>Biological Activity:</b>	Testing in progress		
<b>Predicted N-terminal:</b>	His		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution.		
<b>Storage &amp; Stability:</b>	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. <b>Avoid repeated freeze-thaw cycles.</b>		

## SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)