

## TNFSF12

### Recombinant Human TWEAK

<b>Catalog No.</b>	CRT600B	<b>Quantity:</b>	25 µg
<b>Alternate Names:</b>	Tumor necrosis factor ligand superfamily member 12, APO3 ligand, TNF-related weak inducer of apoptosis, TWEAK		
<b>Description:</b>	TWEAK belongs to the TNF family of ligands and signals through TWEAKR also known as TNFRSF12A. TWEAK is expressed in a variety of tissues, including the adult heart, pancreas, skeletal muscle, small intestine, spleen and peripheral blood lymphocytes. TWEAK has the ability to induce NFκB activation and chemokine secretion, and to exert an apoptotic activity in certain cells, such as HT-29 human adenocarcinoma cells when cultured in the presence of IFN-γ. TWEAK also promotes proliferation and migration of endothelial cells. Recombinant human TWEAK is a soluble 17.0 kDa polypeptide (154 amino acid residues) comprising the TNF homologous region of TWEAK and is generated by proteolytic processing of the full length membrane anchored TWEAK protein.		
<b>UniProt ID:</b>	O43508		
<b>Gene ID:</b>	8742		
<b>Source:</b>	<i>E. coli</i>		
<b>Molecular Weight:</b>	17.0 kDa (154 aa)		
<b>Formulation:</b>	Lyophilized from 10 mM sodium phosphate, pH 7.5 + 0.1 M L-Arg		
<b>Purity:</b>	> 98% as determined by SDS-PAGE and HPLC analyses		
<b>Endotoxin Level:</b>	< 1 EU/µg protein		
<b>Biological Activity:</b>	<p><b>Assay #1:</b> The ED<sub>50</sub> as determined by the dose-dependent stimulation of IL-8 production by Human PBMC is less than 10 ng/ml.</p> <p><b>Assay #2:</b> TWEAK weakly induces the death of HT29 cells when cultured in the presence of IFN-gamma. The ED<sub>50</sub> for this effect is between 30 - 45 ng/ml.</p>		
<b>Amino Acid Sequence:</b>	MKGRKTRARR AIAAHYEVHP RPGQDGAQAG VDGTVSGWEE ARINSSSPLR YNRQIGEFIV TRAGLYLYYC QVHFDEGKAV YLKL DLLVDG VLALRCLEEF SATAASSLGP QLRLCQVSGL LALRPGSSLR IRTLPWAHLK AAPFLTYFGL FQVH		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> First add 10 mM sodium phosphate to the vial to fully solubilize the protein to a concentration of 1.0 mg/ml. After complete solubilization of the protein, it can be further diluted to other aqueous solutions.		
<b>Storage &amp; Stability:</b>	Store lyophilized protein at -20°C to -80°C for up to 1 year. Reconstituted protein is stable for immediate use at 2-8°C or in working aliquots at -20°C to -80°C for longer storage. <b>Avoid repeated freeze-thaw cycles.</b>		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)