

II21

Recombinant Mouse Interleukin-21, Animal Free

Catalog No.	CS322A-AF	Quantity:	2 µg
	CS322B-AF		10 µg
	CS322C-AF		1 mg
	CS322D-AF		100 µg

Alternate Names: IL-21

Description: Interleukin-21 (IL-21) is a member of the common gamma-chain family of cytokines (includes IL-2, IL-4, IL-7, IL-9, IL-13, and IL-15) and is expressed by activated CD4+ T cells. IL-21 has a private receptor known as IL-21R, which is expressed on T, B, and NK cells. Within the B cell lineage, IL-21 regulates IgG1 production and cooperates with IL-4 for the production of multiple Ab classes in vivo. Initial studies have demonstrated that IL-21 has pleiotropic effects on the proliferation, differentiation, and effector functions of B, T, natural killer, and dendritic cells.

Gene ID: 60505

UniProt ID: Q9ES17

Source: *E. coli*

Manufactured without Animal-derived products, in an Animal Free facility.

Molecular Weight: 15.1 kDa (130 aa) monomer

Formulation: Lyophilized from a sterile filtered solution containing 20 mM NaHCO₃, pH 8.5.

Purity: ≥ 95% determined by reducing and non-reducing SDS-PAGE.

Endotoxin Level: ≤ 1 EU/µg by kinetic LAL analysis.

Biological Activity: ED₅₀ ≤ 20 ng/ml, determined by a dose-dependent proliferation of B9 cells.

Specific Activity: ≥ 5.0 x 10⁴ units/mg



Amino Acid Sequence: MHKSSPQGPD RLLIRLRHLI DIVEQLKIYE NDLDPELLSA PQDVKGHCEH
AAFACFQKAK LKPSNPGNNK TFIIDLVAQL RRRLPARRGG KKQKHIKACP
SCDSYEK RTP KEFLERLKW LQKMIHQHLS

Reconstitution: **Centrifuge vial before opening.** When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.

Storage & Stability: Upon receipt, store at -20°C to -80°C for up to 1 year. Upon reconstitution, the preparation is stable for up to one month at 2-8°C. For long term storage, prepare working aliquots and store at -20 to -80°C. For maximal stability, dilute to working aliquots in a 0.1% BSA solution.

Avoid repeated freeze-thaw cycles.





Mouse IL-21 Gel

Figure: 1 ug run under (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Mouse IL-21 is predicted to have a MW of 15.1 kDa.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com