cellsciences.com

Raji
Whole Cell Lysate

Catalog No: PX325A **Size:** 100 µg
PX325B **Size:** 500 µg

Cell Line: Raji (Human Burkitt's lymphoma)

Growth Media: RPMI & 10% NCS (Newborn calf serum)

Cell Lysate Preparation: Raji cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% beta-mercaptoethanol.

Application: Raji cell lysate is ready to load on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

Storage: It is supplied as whole cell lysate at 100 µg at 2 mg/ml in 1 x SDS sample buffer containing 5% beta-mercaptoethanol. Store at -20°C for three months or at –70°C for one year.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Cell Sciences, Inc.
480 Neponset Street
Building 12A
Canton, MA 02021

Toll Free: 888 769-1246
Phone: 781 828-0610
Fax: 781 828-0542

E-mail: info@cellsciences.com
Web Site: www.cellsciences.com